



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence, substituting the values of KN and OK ,

$$\frac{\tan x + \tan u}{1 - \tan x \tan u} = \frac{2 \cot z - \cos y}{\sin y}.$$

Substituting the value of $\tan u$,

$$2 \sin y (\tan z - \cot z) = 2 \tan x \cos y (\cot z + \tan z) - 5 \tan x.$$

$$\text{But } \tan z - \cot z = -2 \tan y \text{ and } \cot z + \tan z = \frac{2}{\cos y}.$$

Hence, substituting and reducing,

$$\tan x = 4 \tan y \sin y,$$

the required relation.


Also solved by PAUL CAPRON and J. W. CLAWSON.

458. Proposed by CLIFFORD N. MILLS, Brookings, S. Dak.

Given edges l , m and n of a parallelopiped and angles a , b and c which the edges make with one another. Show that, if $s = \frac{a + b + c}{2}$, the volume equals

$$2lmn \sqrt{\sin s \sin (s - a) \sin (s - b) \sin (s - c)}.$$

SOLUTION BY FRANK R. MORRIS, Glendale, Cal.



Given the parallelopiped AH with $\angle BAC = a$, $\angle DAC = b$, $\angle BAD = c$, and $AB = l$, $AC = m$, $AD = n$.

The area of the base BD is equal to $ln \sin C$.

From the vertex C drop a perpendicular to the base BD meeting it at E . From E draw perpendiculars to AB and AD meeting the lines at F and G , respectively. CE is the altitude of the parallelopiped and we know that the volume is $V = CE \cdot ln \sin c$. (1).

The triangles AEC , AFC , AGC , AFE and AGE are right triangles. Hence, we have

$$CE^2 = m^2 - AE^2, \quad (2) \quad AF = m \cos a, \quad AG = m \cos b, \quad (3)$$

$$\cos \angle EAF = \frac{AF}{AE}, \quad (4) \quad \text{and} \quad \cos (c - \angle EAF) = \frac{AG}{AE},$$

or

$$\cos c \cos \angle EAF + \sin c \sqrt{1 - \cos^2 \angle EAF} = \frac{AG}{AE}. \quad (5)$$

Eliminating $\cos \angle EAF$ from (4) and (5)

$$\cos c \frac{AF}{AE} + \sin c \sqrt{1 - \frac{AF^2}{AE^2}} = \frac{AG}{AE}.$$

From this equation we get

$$AE^2 = \frac{AG^2 - 2AG \cdot AF \cos c + AF^2}{\sin^2 c}.$$

Substituting the values of AF and AG from (3),

$$AE^2 = \frac{m^2 \cos^2 b - 2m^2 \cos a \cos b \cos c + m^2 \cos^2 a}{\sin^2 c}.$$

Then from (2),

$$\begin{aligned} CE^2 &= \frac{m^2}{\sin^2 c} (\sin^2 c - \cos^2 b + 2 \cos a \cos b \cos c - \cos^2 a) \\ &= \frac{m^2}{\sin^2 c} (\sin^2 c + \cos^2 c - \cos^2 b - \cos^2 c + \cos^2 b \cos^2 c - \cos^2 b \cos^2 c \\ &\quad + 2 \cos a \cos b \cos c - \cos^2 a) \\ &= \frac{m^2}{\sin^2 c} \{ (1 - \cos^2 b - \cos^2 c + \cos^2 b \cos^2 c) - \cos^2 a + 2 \cos a \cos b \cos c - \cos^2 b \cos^2 c \} \\ &= \frac{m^2}{\sin^2 c} \{ (1 - \cos^2 b)(1 - \cos^2 c) - \cos^2 a + 2 \cos a \cos b \cos c - \cos^2 b \cos^2 c \} \\ &= \frac{m^2}{\sin^2 c} \{ \sin^2 b \sin^2 c - \cos^2 a + 2 \cos a \cos b \cos c - \cos^2 b \cos^2 c \} \\ &= \frac{m^2}{\sin^2 c} \{ \sin b \sin c + (\cos a - \cos b \cos c) \} \{ \sin b \sin c - (\cos a - \cos b \cos c) \} \\ &= \frac{m^2}{\sin^2 c} \{ \cos a - \cos(b + c) \} \{ \cos(b - c) - \cos a \} \\ &= \frac{m^2}{\sin^2 c} \left\{ 2 \sin \left(\frac{a + b + c}{2} \right) \sin \left(\frac{b + c - a}{2} \right) \right\} \left\{ \sin \left(\frac{a - b + c}{2} \right) \sin \left(\frac{a + b - c}{2} \right) \right\} \\ &= \frac{4m^2}{\sin^2 c} \sin s \cdot \sin(s - a) \sin(s - b) \sin(s - c), \end{aligned}$$

where

$$s = \frac{a + b + c}{2}.$$

Hence, $CE = \frac{2m}{\sin c} \sqrt{\sin s \sin(s - a) \sin(s - b) \sin(s - c)}$. Substituting this value of CE in (1), we have

$$v = 2lmn \sqrt{\sin s \sin(s - a) \sin(s - b) \sin(s - c)}.$$

Also solved by GEORGE W. HARTWELL, HORACE OLSON, J. A. CAPARO, and J. W. CLAWSON.

CALCULUS.

370. Proposed by PAUL CAPRON, Annapolis, Maryland.

The surface of a right circular cone having the semi-vertical angle α is cut by two planes, which intersect the axis at the same point, one at right angles to the axis, the other making the angle $(90^\circ - \beta)$ with the axis. Show that if the lateral surface of the right cone is S_1 and that of the oblique cone S_2 ,

$$S_2 = \sum_{n=1}^{\infty} T_n, \quad \text{where} \quad T_1 = S_1, \quad T_{n+1} = T_n \times \frac{2n+1}{2n} (\tan \alpha \tan \beta)^2.$$

SOLUTION BY THE PROPOSER.

Let the vertex of the cone be the origin, its axis the z -axis, and let the intersection of the oblique plane with the xy -plane be parallel to the y -axis. Let the two planes cut the z -axis at $(0, 0, h)$, and let the radius of the right section be a . Then,